

1. Physicochemical Property

| Property | Value | Comment |
|------------------|---------|--|
| Molecular Weight | 343.07 | Contain hydrogen atoms. Optimal:100~600 |
| Volume | 327.763 | Van der Waals volume |
| Density | 1.047 | Density = MW / Volume |
| nHA | 6 | Number of hydrogen bond acceptors. Optimal:0~12 |
| nHD | 1 | Number of hydrogen bond donors. Optimal:0~7 |
| nRot | 4 | Number of rotatable bonds. Optimal:0~11 |
| nRing | 3 | Number of rings. Optimal:0~6 |
| MaxRing | 6 | Number of atoms in the biggest ring. Optimal:0~18 |
| nHet | 7 | Number of heteroatoms. Optimal:1~15 |
| fChar | 0 | Formal charge. Optimal:-4 ~4 |
| nRig | 19 | Number of rigid bonds. Optimal:0~30 |
| Flexibility | 0.211 | Flexibility = nRot / nRig |
| Stereo Centers | 0 | Optimal: ≤ 2 |
| TPSA | 80.93 | Topological Polar Surface Area. Optimal:0~140 |
| logS | -4.653 | Log of the aqueous solubility. Optimal: -4~0.5 log mol/L |
| logP | 3.525 | Log of the octanol/water partition coefficient. Optimal: 0~3 |
| logD | 3.383 | logP at physiological pH 7.4. Optimal: 1~3 |

2. Medicinal Chemistry

| Property | Value | Decision | Comment |
|----------|-------|----------|--|
| QED | 0.58 | ● | <ul style="list-style-type: none"> ■ A measure of drug-likeness based on the concept of desirability; ■ Attractive: > 0.67; unattractive: 0.49~0.67; too complex: < 0.34 |
| SAscore | 2.262 | ● | <ul style="list-style-type: none"> ■ Synthetic accessibility score is designed to estimate ease of synthesis of drug-like molecules. ■ SAscore ≥ 6, difficult to synthesize; SAscore <6, easy to synthesize |
| Fsp3 | 0.118 | ● | <ul style="list-style-type: none"> ■ The number of sp³ hybridized carbons / total carbon count, correlating with melting point and solubility. ■ Fsp³ ≥ 0.42 is considered a suitable value. |
| MCE-18 | 18.0 | ● | <ul style="list-style-type: none"> ■ MCE-18 stands for medicinal chemistry evolution. ■ MCE-18 ≥ 45 is considered a suitable value. |

| | | | |
|-----------------|----------|---|---|
| NPscore | -1.491 | - | <p>■ Natural product-likeness score.</p> <p>■ This score is typically in the range from -5 to 5. The higher the score is, the higher the probability is that the molecule is a NP.</p> |
| Lipinski Rule | Accepted | ● | <p>■ $MW \leq 500$; $\log P \leq 5$; $Hacc \leq 10$; $Hdon \leq 5$</p> <p>■ If two properties are out of range, a poor absorption or permeability is possible, one is acceptable.</p> |
| Pfizer Rule | Accepted | ● | <p>$\log P > 3$; $TPSA < 75$</p> <p>Compounds with a high log P (>3) and low TPSA (<75) are likely to be toxic.</p> |
| GSK Rule | Accepted | ● | <p>■ $MW \leq 400$; $\log P \leq 4$</p> <p>■ Compounds satisfying the GSK rule may have a more favorable ADMET profile</p> |
| Golden Triangle | Accepted | ● | <p>■ $200 \leq MW \leq 500$; $-2 \leq \log D \leq 5$</p> <p>■ Compounds satisfying the Golden Triangle rule may have a more favorable ADMET profile.</p> |
| PAINS | 0 alerts | - | Pan Assay Interference Compounds, frequent hitters, Alpha-screen artifacts and reactive compound. |
| ALARM NMR | 1 alerts | - | Thiol reactive compounds. |
| BMS | 0 alerts | - | Undesirable, reactive compounds. |
| Chelator Rule | 0 alerts | - | Chelating compounds. |

3. Absorption

| Property | Value | Decision | Comment |
|---------------------|---------|----------|---|
| Caco-2 Permeability | -4.83 | ● | Optimal: higher than -5.15 Log unit |
| MDCK Permeability | 6.5e-05 | ● | <p>■ low permeability: $< 2 \times 10^{-6}$ cm/s</p> <p>■ medium permeability: $2-20 \times 10^{-6}$ cm/s</p> <p>■ high passive permeability: $> 20 \times 10^{-6}$ cm/s</p> |
| Pgp-inhibitor | 0.207 | ● | <p>■ Category 1: Inhibitor; Category 0: Non-inhibitor;</p> <p>■ The output value is the probability of being Pgp-inhibitor</p> |
| Pgp-substrate | 0.001 | ● | <p>■ Category 1: substrate; Category 0: Non-substrate;</p> <p>■ The output value is the probability of being Pgp-substrate</p> |
| HIA | 0.004 | ● | <p>■ Human Intestinal Absorption</p> <p>■ Category 1: HIA+ (HIA < 30%); Category 0: HIA- (HIA < 30%); The output value is the probability of being HIA+</p> |
| F _{20%} | 0.001 | ● | <p>■ 20% Bioavailability</p> <p>■ Category 1: F_{20%} + (bioavailability < 20%); Category 0: F_{20%} - (bioavailability ≥ 20%); The output value is the probability of being F_{20%} +</p> |

| | | | |
|------------|-----|---|--|
| $F_{30\%}$ | 0.0 | ● | ■ 30% Bioavailability ■ Category 1: $F_{30\%} +$ (bioavailability < 30%); Category 0: $F_{30\%} -$ (bioavailability \geq 30%); The output value is the probability of being $F_{30\%} +$ |
|------------|-----|---|--|

4. Distribution

| Property | Value | Decision | Comment |
|-----------------|--------|----------|--|
| PPB | 99.78% | ● | ■ Plasma Protein Binding ■ Optimal: < 90%. Drugs with high protein-bound may have a low therapeutic index. |
| VD | 0.533 | ● | ■ Volume Distribution ■ Optimal: 0.04-20L/kg |
| BBB Penetration | 0.691 | ● | ■ Blood-Brain Barrier Penetration ■ Category 1: BBB+; Category 0: BBB-; The output value is the probability of being BBB+ |
| Fu | 0.833% | ● | ■ The fraction unbound in plasms ■ Low: <5%; Middle: 5~20%; High: > 20% |

5. Metabolism

| Property | Value | Comment |
|-------------------|-------|--|
| CYP1A2 inhibitor | 0.861 | ■ Category 1: Inhibitor; Category 0: Non-inhibitor; ■ The output value is the probability of being inhibitor. |
| CYP1A2 substrate | 0.622 | ■ Category 1: Substrate; Category 0: Non-substrate; ■ The output value is the probability of being substrate. |
| CYP2C19 inhibitor | 0.938 | ■ Category 1: Inhibitor; Category 0: Non-inhibitor; ■ The output value is the probability of being inhibitor. |
| CYP2C19 substrate | 0.154 | ■ Category 1: Substrate; Category 0: Non-substrate; ■ The output value is the probability of being substrate. |
| CYP2C9 inhibitor | 0.934 | ■ Category 1: Inhibitor; Category 0: Non-inhibitor; ■ The output value is the probability of being inhibitor. |
| CYP2C9 substrate | 0.902 | ■ Category 1: Substrate; Category 0: Non-substrate; ■ The output value is the probability of being substrate. |
| CYP2D6 inhibitor | 0.737 | ■ Category 1: Inhibitor; Category 0: Non-inhibitor; ■ The output value is the probability of being inhibitor. |
| CYP2D6 substrate | 0.768 | ■ Category 1: Substrate; Category 0: Non-substrate; ■ The output value is the probability of being substrate. |
| CYP3A4 inhibitor | 0.912 | ■ Category 1: Inhibitor; Category 0: Non-inhibitor; ■ The output value is the probability of being inhibitor. |
| CYP3A4 substrate | 0.886 | ■ Category 1: Substrate; Category 0: Non-substrate; ■ The output value is the probability of being substrate. |

6. Excretion

| Property | Value | Decision | Comment |
|------------------|-------|----------|---|
| CL | 2.174 | ● | <ul style="list-style-type: none"> ■ Clearance ■ High: >15 mL/min/kg; moderate: 5-15 mL/min/kg; low: <5 mL/min/kg |
| T _{1/2} | 0.22 | - | <ul style="list-style-type: none"> ■ Category 1: long half-life ; Category 0: short half-life; ■ long half-life: >3h; short half-life: <3h ■ The output value is the probability of having long half-life. |

7. Toxicity

| Property | Value | Decision | Comment |
|-------------------------|-------|----------|---|
| hERG Blockers | 0.252 | ● | <ul style="list-style-type: none"> ■ Category 1: active; Category 0: inactive; ■ The output value is the probability of being active. |
| H-HT | 0.405 | ● | <ul style="list-style-type: none"> ■ Human Hepatotoxicity ■ Category 1: H-HT positive(+); Category 0: H-HT negative(-); ■ The output value is the probability of being toxic. |
| DILI | 0.947 | ● | <ul style="list-style-type: none"> ■ Drug Induced Liver Injury. ■ Category 1: drugs with a high risk of DILI; Category 0: drugs with no risk of DILI. The output value is the probability of being toxic. |
| AMES Toxicity | 0.894 | ● | <ul style="list-style-type: none"> ■ Category 1: Ames positive(+); Category 0: Ames negative(-); ■ The output value is the probability of being toxic. |
| Rat Oral Acute Toxicity | 0.09 | ● | <ul style="list-style-type: none"> ■ Category 0: low-toxicity; Category 1: high-toxicity; ■ The output value is the probability of being highly toxic. |
| FDAMDD | 0.175 | ● | <ul style="list-style-type: none"> ■ Maximum Recommended Daily Dose ■ Category 1: FDAMDD (+); Category 0: FDAMDD (-) ■ The output value is the probability of being positive. |
| Skin Sensitization | 0.65 | ● | <ul style="list-style-type: none"> ■ Category 1: Sensitizer; Category 0: Non-sensitizer; ■ The output value is the probability of being sensitizer. |
| Carcinogenicity | 0.645 | ● | <ul style="list-style-type: none"> ■ Category 1: carcinogens; Category 0: non-carcinogens; ■ The output value is the probability of being toxic. |
| Eye Corrosion | 0.003 | ● | <ul style="list-style-type: none"> ■ Category 1: corrosives ; Category 0: noncorrosives ■ The output value is the probability of being corrosives. |
| Eye Irritation | 0.286 | ● | <ul style="list-style-type: none"> ■ Category 1: irritants ; Category 0: nonirritants ■ The output value is the probability of being irritants. |

| | | | |
|----------------------|-------|---|--|
| Respiratory Toxicity | 0.929 | ● | <ul style="list-style-type: none"> ■ Category 1: respiratory toxicants; Category 0: respiratory nontoxicants ■ The output value is the probability of being toxic. |
|----------------------|-------|---|--|

8. Environmental toxicity

| Property | Value | Comment |
|--------------------------|-------|--|
| Bioconcentration Factors | 0.935 | <ul style="list-style-type: none"> ■ Bioconcentration factors are used for considering secondary poisoning potential and assessing risks to human health via the food chain. ■ The unit is $-\log_{10}[(\text{mg/L})/(1000 \cdot \text{MW})]$ |
| IGC ₅₀ | 4.265 | <ul style="list-style-type: none"> ■ Tetrahymena pyriformis 50 percent growth inhibition concentration ■ The unit is $-\log_{10}[(\text{mg/L})/(1000 \cdot \text{MW})]$ |
| LC ₅₀ FM | 4.887 | <ul style="list-style-type: none"> ■ 96-hour fathead minnow 50 percent lethal concentration ■ The unit is $-\log_{10}[(\text{mg/L})/(1000 \cdot \text{MW})]$ |
| LC ₅₀ DM | 4.875 | <ul style="list-style-type: none"> ■ 48-hour daphnia magna 50 percent lethal concentration ■ The unit is $-\log_{10}[(\text{mg/L})/(1000 \cdot \text{MW})]$ |

9. Tox21 pathway

| Property | Value | Decision | Comment |
|---------------|-------|----------|--|
| NR-AR | 0.02 | ● | <ul style="list-style-type: none"> ■ Androgen receptor ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| NR-AR-LBD | 0.01 | ● | <ul style="list-style-type: none"> ■ Androgen receptor ligand-binding domain ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| NR-AhR | 0.768 | ● | <ul style="list-style-type: none"> ■ Aryl hydrocarbon receptor ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| NR-Aromatase | 0.731 | ● | <ul style="list-style-type: none"> ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| NR-ER | 0.589 | ● | <ul style="list-style-type: none"> ■ Estrogen receptor ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| NR-ER-LBD | 0.078 | ● | <ul style="list-style-type: none"> ■ Estrogen receptor ligand-binding domain ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| NR-PPAR-gamma | 0.176 | ● | <ul style="list-style-type: none"> ■ Peroxisome proliferator-activated receptor gamma ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| SR-ARE | 0.495 | ● | <ul style="list-style-type: none"> ■ Antioxidant response element ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| SR-ATAD5 | 0.028 | ● | <ul style="list-style-type: none"> ■ ATPase family AAA domain-containing protein 5 ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |

| | | | |
|--------|-------|---|--|
| SR-HSE | 0.012 | ● | <ul style="list-style-type: none"> ■ Heat shock factor response element ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| SR-MMP | 0.812 | ● | <ul style="list-style-type: none"> ■ Mitochondrial membrane potential ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |
| SR-p53 | 0.616 | ● | <ul style="list-style-type: none"> ■ Category 1: actives ; Category 0: inactives; ■ The output value is the probability of being active. |

10. Toxicophore Rules

| Property | Value | Comment |
|-----------------------------------|----------|---|
| Acute Toxicity Rule | 0 alerts | <ul style="list-style-type: none"> ■ 20 substructures ■ acute toxicity during oral administration |
| Genotoxic Carcinogenicity Rule | 5 alerts | <ul style="list-style-type: none"> ■ 117 substructures ■ carcinogenicity or mutagenicity |
| NonGenotoxic Carcinogenicity Rule | 1 alerts | <ul style="list-style-type: none"> ■ 23 substructures ■ carcinogenicity through nongenotoxic mechanisms |
| Skin Sensitization Rule | 0 alerts | <ul style="list-style-type: none"> ■ 155 substructures ■ skin irritation |
| Aquatic Toxicity Rule | 1 alerts | <ul style="list-style-type: none"> ■ 99 substructures ■ toxicity to liquid(water) |
| NonBiodegradable Rule | 3 alerts | <ul style="list-style-type: none"> ■ 19 substructures ■ non-biodegradable |
| SureChEMBL Rule | 0 alerts | <ul style="list-style-type: none"> ■ 164 substructures ■ MedChem unfriendly status |